

Impact of Prenatal Tobacco Smoke Exposure, as Measured by Midgestation Serum Cotinine Levels, on Fetal Biometry and Umbilical Flow Velocity Waveforms

Jarosław Kalinka, M.D., Ph.D.,¹ Wojciech Hanke,² and Wojciech Sobala²

ABSTRACT

The aim of this prospective cohort study was to evaluate the impact of tobacco smoke exposure, measured by maternal serum concentration of cotinine, on fetal mid-gestation biometric parameters and umbilical artery (UA) qualitative blood flow indices. The study population consisted of 114 healthy women in 20 to 24 weeks gestation who were recruited from the patients of two antenatal care units in Lodz, Poland. Significant negative correlation was found between fetal biparietal diameter (BPD) and serum cotinine concentration. Serum cotinine positively correlated with all blood flow indices under study (systolic/diastolic index [S/D], resistance index, and pulsatility index) after controlling for gestational age, gender, and femur length. The midgestation UA S/D ratio > 3 was found to be a significant risk factor for decreased birthweight. Tobacco smoke exposure is a significant factor inducing increased resistance of umbilical blood flow as measured in 20 to 24 weeks gestation. This could be one of the main mechanisms leading to decreased birthweight observed among infants with prenatal exposure to tobacco smoke.

KEYWORDS: Environmental tobacco smoke exposure, cotinine, uteroplacental circulation

The association between maternal cigarette smoking during pregnancy and the elevated risk for low birthweight, preterm delivery (PD), and intrauterine growth restriction has been well documented in many studies.¹⁻³ According to Kyrklund-Blomberg and Cnattingius,² there is a dose-related impact of smoking on the risk of preterm delivery. The highest impact refers to the risk of spontaneous very preterm births among women smoking at least 10 cigarettes a day (odds ratio, 1.7).

The sidestream smoke (SS), the main constituent of environmental tobacco smoke (ETS), contains similar constituents as does the cigarette smoke inhaled by active smokers, namely tar, nicotine, carbon mono- and dioxide, and benzo(a)pyrene. Moreover, the concentrations of the chemicals found in SS may (even by several times) exceed those in the smoke from tobacco combustion (mainstream smoke).⁴⁻⁸ Many of these substances are known to pass through the placental barrier.^{9,10}

American Journal of Perinatology, Volume 22, 2005. Address for correspondence and reprint requests: Jarosław Kalinka, M.D., Ph.D., Department of Perinatology, Institute of Gynecology and Obstetrics, Medical University of Lodz, ul. Wilenska 37, 94-029 Lodz, Poland. ¹Department of Perinatology, Institute of Gynecology and Obstetrics, Medical University of Lodz, Poland; ²Department of Environmental Epidemiology, Nofer Institute of Occupational Medicine, Lodz, Poland. Copyright © 2005 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662. DOI 10.1055/s-2004-837266. 0735-1631.

Given that the chemical composition of ETS differs quantitatively from the composition of the mainstream smoke, a question arises whether a nonsmoking pregnant woman exposed to ETS toxicants could also be at a higher risk for similar adverse effects on fetal health as is the smoking woman.

The impact of exposure to ETS (passive smoking) on pregnancy outcome has also been investigated in several studies, but the results are often controversial.¹¹⁻¹⁷ An increased risk was found for a small for gestational age (SGA) infant^{12,14} and low birthweight (LBW) in term deliveries.¹⁶ However, negative results^{11,17} or borderline excess risk of LBW were also reported.¹⁸ A relationship between ETS exposure and the rate of PD was postulated by Ahlborg et al¹¹ and confirmed for women with the highest levels of ETS exposure in comprehensive studies of pregnant women from central Poland¹⁵ and the Czech Republic.¹³ The contrasting results might be explained by the differences in the methodology used for ETS exposure assessment (questionnaire versus biomarkers) as well as different study populations, gestational age at the time of ETS exposure assessment, and definitions of the examined endpoints. Klebanoff et al¹⁹ showed that pregnant women accurately reported whether they smoked, but the serum concentration of cotinine, the major metabolite of nicotine, was a better measure of the actual tobacco smoke exposure.

Hayde et al²⁰ observed that maternal cotinine levels correlated with maternal and fetal carboxyhemoglobin (COHb) levels, which suggests that maternal cotinine measurements may be interpreted as the surrogates of fetal COHb.

Few studies addressed the issue of the impact of ETS exposure on fetal biometry and placental vascular resistance.²¹ The mechanism underlying the adverse effect of passive smoking on the course and outcome of pregnancy might be complex. One may presume that a long-term exposure to tobacco smoke constituents, even at low doses, may lead to an impaired fetomaternal blood flow circulation.²² The blood flow velocity changes may be mediated by nicotine, carbon monoxide, catecholamines, or some other factors related to smoking. It has been speculated that smoking causes an increase in the placental vascular resistance both in humans and animals.^{21,23,24}

According to Newnham et al,²⁵ to explain the association between tobacco smoke exposure during pregnancy and fetal growth, there is a need for studies that would be based on prenatal ultrasound measurements rather than the findings of the infant's examination after birth.

The main aim of this prospective cohort study was to evaluate the impact of tobacco smoke exposure, measured by serum concentration of cotinine, on fetal

midgestation biometric parameters and umbilical artery (UA) qualitative blood flow indices.

MATERIALS AND METHODS

Subjects

The study population comprised a group of 129 healthy women in 20 to 24 weeks gestation who were enrolled consecutively at two antenatal care units in Lodz, central Poland, over a 1-year period. Those two units served mostly an indigent population characterized by a high prevalence of smoking and ETS exposure, compared with communities from other regions of central Poland.¹¹ Only the singleton pregnancies were qualified for inclusion in the survey. Women with chronic diseases diagnosed during the first visit were not considered. A standard questionnaire covering medical, socioeconomic, demographic, and constitutional aspects as well as tobacco smoking, including ETS exposure, was administered to every subject and verified based on medical records. Routine ultrasound examination of fetal biometry and Doppler umbilical blood flow assessment was performed along with serum cotinine determination for assessment of tobacco smoke exposure. For five subjects, complete medical records of the newborns were not available. Ten women refused to have serum cotinine measurement ($n=8$) or Doppler ultrasound examination ($n=2$). This prospective cohort study was approved by the Ethical Committee of the Medical University of Lodz, Poland (Decision No. RNN/212/97). Each participant provided a written consent for participation in the study. After ultrasound examination, women who reported smoking or ETS exposure were strongly advised to stop smoking and avoid ETS exposure.

Ultrasound Examination

Ultrasound biometric measurements of fetal biparietal diameter (BPD), abdominal circumference (AC), and femur length (FL) were performed between 20 and 24 weeks of gestation. Each ultrasound measurement was made three times and the mean values were considered for additional analysis. Apart from routine ultrasound, Doppler flow velocimetry in the UA was conducted at the same time. Fetal Doppler waveform examination was made using pulsed Doppler ultrasound after identifying the appropriate vessel by color flow imaging. For all Doppler studies, a 3.5-MHz transducer was used (Hitachi^{Q2} EUB 315C). Recordings were made from the UA by placing the sample volume in the lumen of the artery, away from the placental and fetal cord insertion. Once the satisfactory waveform was imaged, the image was frozen and the blood flow indices (systolic/diastolic index [S/D], resistance index [RI], and pulsatility index [PI]) were calculated.

Serum Cotinine Measurement

Serum cotinine concentration was determined by gas chromatography with mass spectrometry detector to assess ETS exposure during the morning hours of the same day (blood collection at 1200 to 1300 hours). The limit of detection was 0.16 ng/mL, whereas the limit of quantification was 1.25 ng/mL. The half-life of cotinine averages ~17 hours. With intermittent nicotine exposure, which occurs with cigarette smoking or ETS exposure, cotinine levels remain relatively constant throughout the day and remain at near steady-state values.²⁶

In our study, ETS exposure (passive smoking) was diagnosed when the level of serum cotinine ranged between 2 and 14 ng/mL. The subjects regarded as nonsmokers with no ETS exposure had to have serum cotinine levels lower than 2 ng/mL. The subjects were classified as biochemical (active) smokers when their cotinine levels exceeded 14 ng/mL in blood serum.²⁷

Statistical Analysis

The differences between the mean values of quantitative variables were analyzed using linear regression. Multiple linear regressions for BPD, AC, and FL included gestational age at the time of ultrasound examination and fetal gender, whereas those for S/D, RI, and PI also the fetal femur length (FL), were an indicator of fetal size. On the basis of the multiple linear regressions models, the adjusted values of BPD, AC, and FL were calculated for median values in each of the cotinine groups. The differences in the distribution of qualitative variables were evaluated by χ^2 test with Yates correction when necessary. The serum cotinine concentrations were transformed into logarithmic values because they did not meet the criterion of normal distribution.

RESULTS

Study Population Characteristics

The mean age of the examined group was 26.3 years. In the study population, 21% of women were unmarried and 16.1% had primary education. The proportion of women with these characteristics increased with the level of serum cotinine; however, the differences were not statistically significant.

Most of the study group (51.8%) included primiparous women and the mean gestational age was 22.8 weeks. Women with the highest cotinine levels were most frequently primiparous and at a higher gestational age (nonsignificant differences). The proportion of male and female gender of the fetus was well balanced in all of the examined groups, with a slight excess of the male gender in the group with a lower cotinine level (Table 1). As expected, the mean birthweight and gestational age of newborns at delivery was the lowest in the group with highest cotinine levels. The largest proportion of SGA infants (i.e., infants with birthweight below the 10th percentile for a given gestational age, using the national statistics) was found in the group with the highest cotinine levels (9.1%), which, however, was not statistically different from other cotinine groups.

Smoking Status and Serum Cotinine Levels

Tobacco smoking was reported by ~13.2% of the interviewed women (8% admitted to smoking more than five cigarettes a day). Almost half of the nonsmoking women reported daily exposure to ETS. The most common source of ETS exposure was the smoking by family members (30.3%) and smoking at the workplace (13.1%). The results of cotinine determinations indicated that 20% of the 50 persons presumed to be passive

Table 1 Maternal and Pregnancy Characteristics for the Study Groups Classified According to Serum Cotinine Levels

Variable	Serum Cotinine Levels (ng/mL)			p
	< 2.0 (n = 8)	2.0–14 (n = 84)	> 14 (n = 22)	
Maternal characteristics				
Average maternal age, y (SD)	28.0 (4.9)	26.6 (4.9)	24.6 (4.2)	0.146
Average maternal prepregnancy weight ^{Q3} ± SD	57.9 ± 11.1	58.8 ± 9.0	59.3 ± 8.9	0.074
Primary education, n (%)	0 (0.0)	10 (12.2)	8 (36.4)	0.932
Unmarried, n (%)	0 (0.0)	18 (21.4)	6 (27.3)	0.27
Pregnancy characteristics				
Average gestational age at the time of examination (wk) ± SD	21.6 ± 3.9	22.6 ± 3.3	23.7 ± 5.1	0.350
Fetal male gender, n (%)	5 (62.5)	44 (52.4)	8(36.4)	0.31
Primiparous, n (%)	2 (25.0)	44 (52.4)	13 (59.1)	0.25
Pregnancy outcomes				
Average birthweight of newborn (g) ± SD	3280 ± 461	3362 ± 425	3165 ± 568	0.068
Average gestational age at delivery (wk) ± SD	39.3 ± 1.6	39.5 ± 1.6	39.1 ± 1.6	0.137
SGA < 10 percentile, n (%)	0 (0)	4 (4.8)	2 (9.1)	0.57

SD, standard deviation; SGA, small for gestational age.

Table 2 Relationship between Serum Cotinine Level and the Mean Crude and Adjusted Values of Fetal Biometry Assessed between 20 and 24 Weeks of Gestation

Serum Cotinine Level (ng/mL)	n	Fetal Biometry in 20–24 wk of Gestation					
		Crude Values (Mean ± SD)			Adjusted Values*		
		BPD	AC	FL	BPD	AC	FL
< 2.0	8	54.9 ± 10.2	165.5 ± 38.9	37.5 ± 38.9	50.2	152.7	34.0
2.0–14	84	57.0 ± 10.1	180.1 ± 36.4	39.6 ± 8.1	49.7	152.9	33.6
> 14	22	59.0 ± 13.1	190.0 ± 49.8	41.6 ± 11.5	48.6	153.2	32.9
<i>p</i>	0.611	0.309	0.481	0.052	0.681	0.133	

SD, standard deviation; BPD, fetal biparietal diameter; AC, abdominal circumference; FL, femur length.

*Adjusted BPD, AC, and FL values were calculated for the mean value of serum cotinine concentration in each group, 20 weeks of gestation, and female fetus gender, using linear regression model.

smokers were actually actively involved in smoking. On other hand, in the group of 49 women reporting to be nonsmokers not exposed to ETS, the cotinine measurements confirmed this status only in six (12.2%) cases, whereas 43 women (87.8%) appeared to be passive smokers.

Ultrasound Fetal Biometric Measurement and Serum Cotinine Levels

No significant differences were found among the three examined groups (classification based on cotinine level) with respect to the mean BPD, AC, and FL values (Table 2). This could be explained by a slightly older gestational age in the group with the highest cotinine levels. However, after controlling for pregnancy duration and fetal gender in the linear model, a significant negative correlation was found between cotinine concentration and fetal BPD. No correlation of this kind was observed for the AC or FL parameters. The values of respective fetal biometric parameters, calculated using the linear regression model, are listed in Table 2.

Qualitative Indices of Umbilical Blood Flow and Serum Cotinine Levels

Serum cotinine level positively correlated with all blood flow indices under study. The highest mean values were

noted among women classified as active smokers according to their serum cotinine level, and the lowest mean values were among nonsmokers without ETS exposure, with the values for passive smokers in between (Table 3). However, only the differences in the mean values of S/D and PI reached statistical significance.

A linear regression model was constructed to evaluate the relationship between serum cotinine levels and umbilical blood flow indices (Table 3). In the model we also included the fetal FL and fetal gender as the confounding variables. Our previous studies (the findings not shown) indicated that the longer FL, the lower the value of the umbilical artery S/D ratio. On the other hand, the mean value for this index was higher among female fetuses compared with the male fetuses. Cotinine level was found to be a significant predictor for all of the examined UA blood flow indices. On the basis of the linear regression model, adjusted values of three umbilical blood flow indices were calculated for the value of median cotinine concentration in each cotinine group (Table 3).

S/D Index and Infant Birthweight

To evaluate the prognostic value of S/D index, a linear regression model was developed, with birthweight as the dependant variable, and pregnancy duration, S/D, and infant's gender as the independent variables. An increase

Table 3 Serum Cotinine Level Measured between 20 and 24 Weeks of Pregnancy and the Mean Crude and Adjusted Values of Umbilical Artery Blood Flow Indices

Serum Cotinine Level (ng/mL)	n	Umbilical Artery Blood Flow in 20–24 wk of Gestation					
		Crude Values (Mean ± SD)			Adjusted Values*		
		S/D	RI	PI	S/D	RI	PI
< 2.0	8	2.86 ± 0.28	0.65 ± 0.04	1.07 ± 0.09	2.93	0.66	1.15
2.0–14	84	3.07 ± 0.54	0.67 ± 0.06	1.17 ± 0.17	3.18	0.67	1.20
> 14	22	3.53 ± 0.97	0.68 ± 0.11	1.25 ± 0.22	3.66	0.70	1.30
<i>p</i>		0.006	0.496	0.038	0.003	0.004	0.004

SD, standard deviation; S/D, systolic/diastolic index; RI, resistance index; PI, pulsatility index.

*Adjusted S/D, RI, and PI values were calculated for the mean serum cotinine level in each cotinine group, 20 weeks of gestation, male fetus gender, and femur length = 39.9, using linear regression model.

Table 4 Relationship between Umbilical Artery S/D Ratio Assessed in 20–24 Weeks of Gestation and Infants' Birthweight: Crude Values and Mean Adjusted for Pregnancy Duration

S/D Value	Number of Subjects Examined	Birthweight (g)		
		Crude Value		Mean Adjusted for Pregnancy Duration
		Mean	SD	
1–2.5	10	3346	465	3371
2.5–3.0	43	3464	456	3484
3.0–3.5	35	3343	362	3438
3.5+	26	2984	462	3128
<i>p</i>		0.001		0.01

S/D, systolic/diastolic index.

in the umbilical artery S/D ratio > 3.0 in 20 to 24 weeks of pregnancy negatively correlated with infant birthweight (Table 4). When the birthweight values obtained for each of the S/D values were adjusted for pregnancy duration, a 243-g difference was found between the lowest and highest S/D group.

DISCUSSION

The findings of this study demonstrated a significant increase in the umbilical artery S/D velocity ratio^{Q4} with increasing serum cotinine levels, both for the active and passive smokers. It is noteworthy that as the serum cotinine level increases, all of the qualitative indices of umbilical blood flow increase, which suggests an increasing vascular resistance in the placental circulation both among active and passive smokers.

Newnham et al²⁵ observed a decreased fetal BPD diameter in the pregnancies of smokers, with a maximal effect at 24 weeks gestation and only among male fetuses. This effect was not associated with an altered head circumference after birth. Our results indicate that the mean fetal BPD measurements are lower, not only in the group of smokers, but also among women exposed to ETS as measured in 20 to 24 weeks gestation.

Findings consistent with the results of our study were reported by Lymeropoulou et al,²¹ who showed a significant increase in uteroplacental vascular resistance related to increased cotinine levels. Kimya et al²⁸ investigated an acute effect of smoking a single, standard 100-mm, filtered cigarette on uterine artery and UA blood velocity waveforms among pregnant women at the mean gestational age of 28 weeks. The results suggest that there was no significant change in the uterine artery and UA blood velocity waveform indices that could be attributed to the acute effect of smoking, but all uterine artery indices and S/D ratio in UA were statistically higher in the smoking women compared with nonsmokers, both before and after smoking. Newnham et al²⁵

observed no differences in S/D ratios in UA between smokers and nonsmokers, which would indicate that the effects of smoking on the placental vascular resistance were periodic rather than continuous. This is contradictory to the results of our study. Our analysis revealed that either active or passive smoking might have an adverse effect on UA blood flow as early as in midgestation, which implies that the impact of smoking might begin in early pregnancy. According to Morrow et al,²⁹ the smoking of one cigarette has no significant impact on vascular resistance of the uterine artery (S/D), but is connected with a highly significant increase in S/D ratio in UA as measured between 36 and 41 weeks of gestation. This observation would suggest that active smoking might induce a direct increase in vascular resistance of the placenta on the fetal side.

The results of the present study indicate that exposure to tobacco smoke in pregnancy may induce a direct increase in the vascular resistance of the placenta as early as at midgestation, and this pathomechanism may contribute to the decreased birthweight of the newborns. The analysis also revealed that an increase in the umbilical artery S/D ratio > 3.0 in 20 to 24 weeks gestation significantly correlated with the decreased birthweight.

The negative impact on fetal growth could be explained by several different mechanisms. By changing the morphology and function of the placenta,³⁰ smoking in pregnancy may lead to fetal hypoxia through placental vasoconstriction. A reduction in uterine blood flow has been demonstrated in animal and human models.^{23,24} In the study conducted by Clark and Irion,²³ pregnant ewes received maternal intravenous infusions of nicotine, which produced an increase in fetal blood pressure, a decrease in fetal heart rate, and a decrease in umbilical blood flow, with insignificantly altered S/D ratios. The differences in the placental enzyme activity between active and passive smokers, to some extent, could also explain the adverse effects of relatively low exposure levels observed among passive smokers. According to Remmer,⁶ in active smokers, the induction of placental enzymes protects the fetus, whereas the small amount of smoke in passive smokers is unable to induce the enzymes and the smoke remains undetoxified^{Q5}. The results obtained by Jauniaux et al³¹ indicate that higher cotinine levels could be found in fetal fluids and serum than in maternal serum, and that cotinine accumulated in fetal compartments as early as 7 weeks gestation in both active and passive smokers. As revealed by the findings of Steuerer et al,³² smoking increases the consumption of vitamin E, hence there is an overproduction of peroxides and a reduction in prostacyclin production, which may lead to a reduced perfusion of the placenta.

The results of this study indicate that intrauterine growth could be negatively affected by tobacco smoke exposure during pregnancy. Doppler examination should

Q4

Q5

be regarded as an additional diagnostic tool in intrauterine fetal evaluation, and cannot, in itself, be considered a determinant of future fetal growth and development. However, the results of UA Doppler examination in early pregnancy can be used for the assessment of the risk of fetal hypoxia in the populations exposed to specific harmful agents.

CONCLUSIONS

1. The findings of this study suggest that the impaired umbilical blood flow among actively and passively smoking pregnant women begins early in pregnancy and could be one of the mechanisms through which tobacco smoke exposure can affect fetal growth.
2. Prevention measures need to be undertaken to encourage pregnant women to stop smoking and avoid passive exposure to tobacco smoke during pregnancy. Educational activities should be provided to the general population to increase social awareness of the adverse effects, not only of active but also of passive smoking, during pregnancy.

ACKNOWLEDGMENTS

This study was supported by Grant 0654/ PO5/97/12 by the Polish Committee for Scientific Research.

REFERENCES

1. Eskenazi B, Prehn AW, Christianson RE. Passive and active maternal smoking as measured by serum cotinine: the effect on birthweight. *Am J Public Health* 1995;85:395–398
2. Kyrklund-Blomberg NB, Cnattingius S. Preterm birth and maternal smoking: risk related to gestational age and onset of delivery. *Am J Obstet Gynecol* 1998;179:1051–1055
3. Windham GC, Hopkins B, Fenster L, Swan SH. Prenatal active or passive tobacco smoke exposure and the risk of preterm delivery or low birth weight. *Epidemiology* 2000;11:427–433
4. Adams JD, O'Mara-Adams KJ, Hoffmann D. Toxic and carcinogenic agents in undiluted mainstream smoke and sidestream smoke of different types of cigarettes. *Carcinogenesis* 1987;8:729–731
5. Iwase A, Aiba M, Kira S. Respiratory nicotine absorption in non-smoking females during passive smoking. *Int Arch Occup Environ Health* 1991;63:139–143
6. Remmer H. Passively inhaled tobacco smoke: a challenge to toxicology and preventive medicine. *Arch Toxicol* 1987;61:89–104
7. U.S Department of Health, Education, and Welfare. Smoking and health: a report of the Surgeon General. Washington, DC: US GPO; 1979
8. US Environmental Protection Agency (EPA), Department of Health and Human Services. Respiratory health effects of passive smoking: lung cancer and other disorders. NIH publication: Smoking and tobacco control monograph 4. Washington, DC: EPA, Department of Health and Human Services; 1993:93–3605^{Q6}
9. Andersen BD. Cotinine in amniotic fluids from passive smokers. *Lancet* 1982;1:791–31^{Q7}
10. Jordanov JS. Cotinine concentrations in amniotic fluid and urine of smoking, passive smoking and non-smoking pregnant women at term and in the urine of their neonates on 1st day of life. *Eur J Pediatr* 1990;149:734–737
11. Ahlberg G, Bodin L. Tobacco smoke exposure and pregnancy outcome among working women. *Am J Epidemiol* 1991;133:338–347
12. Dejin-Karlsson E, Hanson BS, Ostergren PO, Sjoberg NO, Marsal K. Does passive smoking in early pregnancy increase the risk of small-for-gestational-age infants? *Am J Public Health* 1998;88:1523–1527
13. Dejmeck J, Solansky I, Podrazilova K, Sram RJ. The exposure of nonsmoking and smoking mothers to environmental tobacco smoke during different gestational phases of fetal growth. *Environ Health Perspect* 2002;110:601–606
14. Fortier I, Marcoux S, Brisson J. Passive smoking during pregnancy and risk of delivering a small-for-gestational-age infant. *Am J Epidemiol* 1994;139:294–310
15. Hanke W, Kalinka J, Florek E, Sobala W. Passive smoking and pregnancy outcome in Central Poland. *Hum Exp Toxicol* 1999;18:265–271
16. Martin TR, Bracken MB. Association of low birth weight with passive smoke exposure in pregnancy. *Am J Epidemiol* 1986;124:633–642
17. Ogawa H, Tominaga S, Hori K, Noguchi K, Kanou I, Matsubara M. Passive smoking by pregnant women and fetal growth. *J Epidemiol Community Health* 1991;45:164–168
18. Jedrychowski W. Effects of Poor Air Quality on the Health of Krakow Children, 1st ed. Kraków, Poland: Jagiellonian University Collegium Medicum; 1998
19. Klebanoff MA, Levine RJ, Clemens JD, DerSimonian R, Wilkins DG. Serum cotinine concentration and self-reported smoking during pregnancy. *Am J Epidemiol* 1998;148:259–262
20. Hayde M, Bernaschek G, Stevenson DK, Knight GJ, Haddow JE, Widness JA. Antepartum fetal and maternal carboxyhemoglobin and cotinine levels among cigarette smokers. *Acta Paediatr* 1999;88:327–331
21. Lymperopoulou A, Hainaut F, Crimail P, Durand JL, Locatelli C, Maison C. Tabac et grossesse: recherche d'une correlation cotininemie et Doppler. *J Gynecol Obstet Biol Reprod (Paris)* 1996;25:824–827
22. Castro LC, Allen R, Ogunyemi D, Roll K, Platt LD. Cigarette smoking during pregnancy: acute effects on uterine flow velocity waveforms. *Obstet Gynecol* 1993;81:551–555
23. Clark KE, Irion GL. Fetal hemodynamics response to maternal intravenous nicotine administration. *Am J Obstet Gynecol* 1992;167:1624–1631
24. Suzuki K, Minei LJ, Johnson EE. Effect of nicotine upon uterine blood flow in the pregnant rhesus monkey. *Am J Obstet Gynecol* 1980;136:1009–1013
25. Newnham JP, Patterson L, James I, Reid SE. Effects of maternal cigarette smoking on ultrasonic measurements of fetal growth and on Doppler flow velocity waveforms. *Early Hum Dev* 1990;24:23–36
26. Benowitz NL. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol Rev* 1996;18:188–204

Q6

Q7

27. Cummings SR, Richard RJ. Optimum cutoff points for biochemical validation of smoking status. *Am J Public Health* 1988;78:574-575
28. Kimya Y, Cengiz C, Ozan H, Kolsal N. Acute effects of maternal smoking on uterine and umbilical artery blood velocity waveforms. *J Matern Fetal Investig* 1998;8:79-81
29. Morrow RJ, Ritchie JW, Bull SB. Maternal cigarette smoking: the effects on umbilical and uterine blood flow velocity. *Am J Obstet Gynecol* 1988;159:1069-1071
30. Salafia C, Schiverick K. Cigarette smoking and pregnancy, II: vascular effects. *Placenta* 1999;20:273-279
31. Jauniaux E, Gulbis B, Acharya G, Thiry P, Rodeck C. Maternal tobacco exposure and cotinine levels in fetal fluids in the first half of pregnancy. *Obstet Gynecol* 1999;93:25-29
32. Steuerer A, Rosenbaum P, Heller WD, et al. Einfluss des Rauchverhaltens und der antioxidativen Vitaminkonzentrationen von Schwangeren auf das Geburtsgewicht von Neugeborenen. *Z Geburtshilfe Neonatol* 1999;203:110-114



THIEME

Author Query Form (AJP/40200)

Special Instructions: Author please write responses to queries directly on proofs and then return back.

Q1: Au: The length of the running head was reduced to conserve space.

Q2: Au: Please supply city and country location (city and state if U.S.) for Hitachi.

Q3: Au: Is unit of measure for weight kg?

Q4: Au: Correct with SD as changed to S/D to match other text?

Q5: Au: Is “undetoxified” standard jargon? Can it be changed to “toxic” for clarity?

Q6: Au: Is this a page range or a publication number? Please verify.

Q7: Au: Please verify the page numbers (791-31) (in reference 9 Andersen, 1982).



THIEME

Instructions to Contributors

Dear Contributor:

Enclosed in this document please find the page proofs, copyright transfer agreement (CTA), and offprint order form for your article in the *American Journal of Perinatology*, Volume 22, 2005. Please print this document and complete and return the CTA and offprint form, along with corrected proofs, within 72 hours.

- 1) Please read proofs carefully for **typographical** and **factual** errors only; mark corrections in the margins of the proofs in pencil. **Answer (on the proofs) all editor's queries written in the margins of the proofs.** Check references for accuracy. **Please check on the bottom of the 1st page of your article that your titles and affiliations are correct.** Avoid elective changes, as these are costly and time consuming and will be made at the publisher's discretion.
- 2) Please pay particular attention to the proper placement of figures, tables, and legends. Please provide copies of any formal letters of permission that you have obtained.
- 3) **Please return the corrected proofs, signed copyright transfer agreement, and your offprint order form, with the black and white or color prints of figures, if you received any.**
- 4) As a contributor to this journal you will receive one copy of the journal, at no charge.
 - If you wish to order offprints, **please circle the quantity required** (left column) **and the number of pages in your article.** If you wish to order additional copies of the journal please enter the number of copies on the indicated line.
 - If you do not want to order offprints or journals simply put a slash through the form, **but please return the form.**

Please send all materials back via overnight mail, within 72 hours of receipt, to:

David Stewart, Production Editor
Thieme Medical Publishers
333 Seventh Avenue
New York, NY 10001
Tel: 212-584-4693 Fax: 212-947-1112
E-mail: dstewart@thieme.com

Please do not return your materials to the editor, or the compositor.

Please note: Due to a tight schedule, if the publisher does not receive the return of your article within 7 days of the mail date (from the compositor), the publisher reserves the right to proceed with publication without author changes. Such proofs will be proofread by the editor and the publisher.

Thank you for your contribution to this journal.
David Stewart, Production Editor, Journal Production Department
Thieme Medical Publishers, Inc.

Thieme Medical Publishers, Inc. (the "Publisher") will be pleased to publish your article (the "Work") entitled _____ in the *American Journal of Perinatology*, Volume 22, 2005.

The undersigned Author(s) hereby assigns to the Publisher all rights to the Work of any kind, including those rights protected by the United States Copyright laws.

The Author(s) will be given permission by the Publisher, upon written request, to use all or part of the Work for scholarly or academic purposes, provided lawful copyright notice is given.

If the Work, subsequent to publication, cannot be reproduced and delivered to the Author(s) by the publisher within 60 days of a written request, the Author(s) is given permission to reprint the Work without further request.

The Publisher may grant third parties permission to reproduce all or part of the Work. The Author(s) will be notified as a matter of courtesy, not as a matter of contract. Lawful notice of copyright always will be given.

Check appropriate box below and affix signature.

I Sign for and accept responsibility for transferring copyright of this article to Thieme Medical Publishers, Inc. on behalf of any and all authors.

Author's full name, degrees, professional title, affiliation, and complete address:

Author's printed name, degrees

Professional title

Complete professional address

Author's signature

Date

I prepared this article as part of my official duties as an employee of the United States Federal Government. Therefore, I am unable to transfer rights to Thieme Medical Publishers, Inc.

Author's signature

Date

Contributor Offprint Prices and Order Form for the American Journal of Perinatology
(Effective January 1999)

Please circle the cost of the quantity/page count you require (orders must be in increments of 100)

PAGES IN ARTICLE/COST								
Quantity	1 to 4	5 to 8	9 to 12	13 to 16	17 to 20	21 to 24	25 to 28	29 to 32
100	\$173	\$268	\$363	\$505	\$600	\$696	\$814	\$945
200	\$232	\$375	\$529	\$707	\$862	\$1,004	\$1,171	\$1,349
300	\$292	\$470	\$684	\$909	\$1,123	\$1,301	\$1,527	\$1,741
400	\$351	\$577	\$850	\$1,099	\$1,385	\$1,533	\$1,884	\$2,133
500	\$410	\$672	\$1,004	\$1,301	\$1,646	\$1,919	\$2,240	\$2,537
1000	\$707	\$1,183	\$1,812	\$2,299	\$2,941	\$3,440	\$4,034	\$4,509

Volume No: _____ Page Range (of your article): _____

Article Title: _____

MC/Visa/AmEx No: _____ Exp. Date: _____

Signature: _____

Name: _____

Address: _____

City/State/Zip/Country: _____

Number of **additional** copies of the journal, at the discounted rate of \$20.00 each: _____

Notes

1. The above costs are valid only for orders received before the publication of the journal. Please return the completed form, even if your institution intends to send a P.O. (The P.O. may sometimes be supplied after the journal has been printed.)
2. Orders from outside the United States must be accompanied by payment.
3. Prices are F.O.B. from binder. **A shipping charge will be added to the above costs.**
4. Reprints are printed on the same coated paper as the journal and side-stapled.
5. For larger quantities or late orders, please contact Tyge Burgess at: Phone: (212) 584-4662
Fax (212) 947-1112
E-mail: tburgess@thieme.com